

R.R. Gadyl'shin¹

ON LOCAL PERTURBATIONS OF SCHRÖDINGER OPERATOR ON PLANE

We obtain necessary and sufficient conditions for emerging of small eigenvalue for Schrödinger operator on plane under local operator perturbations. In the case the eigenvalue emerges we construct its asymptotics. The examples are given.

1. INTRODUCTION

The questions of existence of bound states and asymptotics for associated eigenvalues (if they exist) for Schrödinger operator are investigated in [1]–[10]. In [11] in order to study the case of small perturbation of Schrödinger operator in the axis originated by sufficient arbitrary localized second-order operator it was used an approach differing from ones employed in [1]–[10]. This approach gave a simple explanation of "non-regular" (non-obligatory) emerging of eigenvalues under, obviously, regular perturbation; the essence of this approach is as follows. Instead of spectral parameter λ they introduced more natural frequency one k related with spectral parameter by equality $\lambda = -k^2$, where k belonged to complex half-plane $\operatorname{Re} k > 0$. The solutions of both non-perturbed and perturbed equation were continued into complex plane by frequency parameter. Under such continuation the solution of non-perturbed equation had a pole at zero moving under perturbation and the residue at this pole (both for perturbed and non-perturbed problems) was a solution of corresponding homogeneous equation. For non-perturbed problem this residue is a constant considered as exponent in a power $-kt$ with $k = 0$. The side the pole moved to determined the exponential increasing or decreasing of residue for perturbed problem. As a result, if pole moved into half-plane $\mathbb{C}_+ \stackrel{\text{def}}{=} \{k : \operatorname{Re} k > 0\}$, the eigenvalue had emerged, and it had not in the case of shift into the half-plane $\operatorname{Re} k \leq 0$. Direction of this shift depended on the operator of perturbation.

For the small perturbation of Schrödinger operator on plane considered in this work and carried out by localized second-order operator, the situation changes noticeably, because here the solution of non-perturbed equation has a logarithmic branch point instead of pole at $k = 0$, and, therefore, is analytically continued from half-plane into complex plane with a cut along negative real semiaxis. None the less, following the scheme proposed in [11], it is possible to get the necessary and sufficient conditions under those the solution of perturbed equation has a small pole in a right half-plane (or, by other words, a small eigenvalue emerge) and to calculate the asymptotics for it.

¹Bashkir State Pedagogical University, Ufa, Russia. E-mail: gadylshin@bspu.ru

The construction of proof, on the one hand, being close to [11], and, on the other hand, having its some differences (regarding to the "replacement" of pole to logarithmic singularity at non-perturbed equation), it is convenient to keep where possible the notations and style of exposition of work [11].

The structure of the work is the following one. In the second section we give a formulation of the main statement (Theorem 1) and its corollaries. The third section is devoted to the proof of the main statement. In concluding fourth section we show some examples illustrating the main statement.

2. FORMULATION OF THE MAIN RESULT

Hereafter, $W_{2,loc}^j(\mathbb{R}^2)$ is a set of functions defined on $\mathbb{R}^2$ whose restrictions into each bounded domain $D \subset \mathbb{R}^2$ belong to $W_2^j(D)$, $\|\bullet\|_G$ and $\|\bullet\|_{j,G}$ are norms in $L_2(G)$ and $W_2^j(G)$, respectively. Next, let Q be an arbitrary bounded domain in $\mathbb{R}^2$, $L_2(\mathbb{R}^2; Q)$ be a subset of functions from $L_2(\mathbb{R}^2)$ having support in $\overline{Q}$, $\mathcal{L}_\varepsilon$ be a linear operator mapping $W_{2,loc}^j(\mathbb{R}^2)$ into $L_2(\mathbb{R}^2; Q)$ such that $\|\mathcal{L}_\varepsilon[u]\|_Q \leq C(\mathcal{L}) \|u\|_{2,Q}$, where constant $C(\mathcal{L})$ is independent on ε , $0 < \varepsilon \ll 1$,

$$\langle g \rangle = \int_{\mathbb{R}^2} g dx, \quad H_0 = -\Delta, \quad H_\varepsilon = -(\Delta + \varepsilon \mathcal{L}_\varepsilon).$$

Hereinafter, $x = (x_1, x_2)$, $y = (y_1, y_2)$. We define linear operators $A(k) : L_2(\mathbb{R}^2; Q) \rightarrow W_{2,loc}^2(\mathbb{R}^2)$ and $T_\varepsilon^{(0)}(k) : L_2(\mathbb{R}^2; Q) \rightarrow L_2(\mathbb{R}^2; Q)$ in a following way:

$$A(k)g = -\frac{1}{2\pi} \int_{\mathbb{R}^2} \mathcal{K}_0(k|x-y|)g(y) dy,$$

$$T_\varepsilon^{(0)}(k)g = \mathcal{L}_\varepsilon[A(k)g] - \frac{\langle g \rangle \mathcal{L}_\varepsilon[1]}{2\pi} (\ln k + \mathbf{C} - \ln 2),$$

where $\mathcal{K}_0$ is a Basse function of zero order (i.e., $\mathcal{K}_0(z) = \frac{i\pi}{2} \mathcal{H}_0^{(1)}(iz)$, where $\mathcal{H}_0^{(1)}$ is the Hankel function of first kind and zero order), $\mathbf{C}$ is the Euler constant. We denote by $\mathcal{B}(X, Y)$ the Banach space of linear bounded operators from Banach space X into Banach space Y , $\mathcal{B}(X) \stackrel{def}{=} \mathcal{B}(X, X)$, and we will employ the symbol $\mathcal{B}^h(X, Y)$ ($\mathcal{B}^h(X)$) for the set of holomorphic operator-valued functions with values in $\mathcal{B}(X, Y)$ (in $\mathcal{B}(X)$). Let I be the identity operator and $S(t)$ be a circle of radius t with center at zero in $\mathbb{C}$ and let $S_\pm(t) = S(t) \setminus \mathbb{R}^\pm$. Hereinafter $\mathbb{R}^-$ ($\mathbb{R}^+$) is a non-positive (non-negative) real semiaxis in a complex plane. Since

$$\mathcal{K}_0(z) = -\ln \frac{z}{2} \sum_{j=0}^{\infty} \frac{z^{2j}}{4^j (j!)^2} + \sum_{j=0}^{\infty} \frac{z^{2j} \psi(j+1)}{4^j (j!)^2},$$

where $\psi(z)$ is a logarithmic derivation of Gamma function (see, for instance, [12], [13]), the definition of $A(k)$ and $T_\varepsilon^{(0)}(k)$ immediately implies

Lemma 1. *For each $R > 0$ there exists $\varepsilon_0(R) > 0$ such that for each fixed $\varepsilon < \varepsilon_0(R)$*

- 1) *there exists $B_\varepsilon(k) \stackrel{\text{def}}{=} (I + \varepsilon T_\varepsilon^{(0)}(k))^{-1} \in \mathcal{B}^h(L_2(\mathbb{R}^2; Q))$ as $k \in S_-(R)$, and also, the uniform convergence $B_\varepsilon(k) \xrightarrow{\varepsilon \rightarrow 0} I$ takes place;*
- 2) *there exists at most one (for each fixed ε) solution $k_\varepsilon \in S_-(R)$ of equation*

$$1 + \varepsilon \frac{1}{2\pi} \langle B_\varepsilon(k) \mathcal{L}_\varepsilon[1] \rangle (\ln k + \mathbf{C} - \ln 2) = 0, \quad (2.1)$$

moreover, if it exists, then

$$k_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} 0. \quad (2.2)$$

Remark 1. The statement of item 2) does not exclude the situation, when for some values ε the solution $k_\varepsilon \in S_-(R)$ of equation (2.1) exists, and for others it doesn't.

We will say the operator $\mathcal{L}_\varepsilon$ to be real-valued if $\text{Im} \langle \overline{g} \mathcal{L}_\varepsilon[g] \rangle = 0$ for each function $g \in W_{2,loc}^2(\mathbb{R}^2)$. Let us denote $\Pi_s(t) = \{k : |\text{Im } k| < sC(\mathcal{L}), \text{Re } k > t\}$. The main contents of this work is the following statement whose proof next section is devoted to.

Theorem 1. *For $\varepsilon \rightarrow 0$ there exists $t(\varepsilon) \rightarrow \infty$, such that in $\mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$ for each fixed ε there is at most one eigenvalue λ_ε of operator H_ε . The criteria of existence of this eigenvalue is an existence of solution $k_\varepsilon \in \mathbb{C}_+$ for the equation (2.1), moreover,*

$$\lambda_\varepsilon = -k_\varepsilon^2, \quad (2.3)$$

and associated unique eigenfunction ϕ_ε is of the form

$$\phi_\varepsilon = A(k_\varepsilon) B_\varepsilon(k_\varepsilon) \mathcal{L}_\varepsilon[1]. \quad (2.4)$$

If, in addition, the operator $\mathcal{L}_\varepsilon$ is real, then the statement of the theorem is true outside the interval $(t(\varepsilon), \infty)$ of real axis.

The assertions (2.1), (2.2) and statement of Theorem 1 yield

Corollary. *If $\mathcal{L}_\varepsilon[1] = 0$, then there is no $\lambda_\varepsilon \notin \mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$.*

If there is no $\lambda_\varepsilon \notin \mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$, then $\lambda_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.

On the one hand, Theorem 1 give neither necessary nor sufficient conditions for operator $\mathcal{L}_\varepsilon$ under those perturbed operator H_ε has small eigenvalues (except necessary condition given in Corollary), and, moreover, this theorem provides no asymptotics for these eigenvalues if they emerge. On the other hand, solving equation (2.1), one can easily deduce these conditions and asymptotics. Indeed, bearing in mind the definition of operators $B_\varepsilon(k)$, $T_\varepsilon^{(0)}(k)$, $A(k)$ and

$$(\Delta^{-1}g)(x) \stackrel{\text{def}}{=} \frac{1}{2\pi} \int_{\mathbb{R}^2} \ln|x-y| g(y) dy,$$

one obtain the correctness of the following statement.

Lemma 2. *If there exists a solution $k_\varepsilon \in S_-(R)$ of equation (2.1), then*

$$k_\varepsilon = \exp(-M(\varepsilon)), \quad (2.5)$$

where $M(\varepsilon)$ has an asymptotics determined by equality

$$M(\varepsilon) = \frac{2\pi}{\varepsilon \sum_{j=0}^{\infty} (-\varepsilon)^j \left\langle (\mathcal{L}_\varepsilon \circ \Delta^{-1})^j \mathcal{L}_\varepsilon[1] \right\rangle} + \mathbf{C} - \ln 2 \quad (2.6)$$

as $\varepsilon \rightarrow 0$, and this asymptotics holds on power precision.

If for the function $\widetilde{M}(\varepsilon)$ equalling to right hand of (2.6) there exists $\alpha > 0$ such that for $\varepsilon \rightarrow 0$ the inequalities

$$\operatorname{Re} \widetilde{M}(\varepsilon) > \alpha \varepsilon^\alpha, \quad \left| \operatorname{Im} \widetilde{M}(\varepsilon) \right| < \frac{\pi}{2} - \alpha \varepsilon^\alpha, \quad (2.7)$$

hold, then for each $R > 0$ there exists $\varepsilon_0(R) > 0$ such that for $\varepsilon < \varepsilon_0(R)$ there exist solutions $k_\varepsilon \in S^+(R) \stackrel{\text{def}}{=} S(R) \cap \mathbb{C}_+$ of equation (2.1).

If for the function $\widetilde{M}(\varepsilon)$ equalling to right hand of (2.6) there exists $\alpha > 0$ such that for $\varepsilon \rightarrow 0$ the inequalities

$$\operatorname{Re} \widetilde{M}(\varepsilon) < -\alpha \varepsilon^\alpha \quad (2.8)$$

or

$$\left| \operatorname{Im} \widetilde{M}(\varepsilon) \right| > \frac{\pi}{2} + \alpha \varepsilon^\alpha, \quad (2.9)$$

hold, then for each $R > 0$ there exists $\varepsilon_0(R) > 0$ such that for $\varepsilon < \varepsilon_0(R)$ there is no solutions $k_\varepsilon \in S^+(R)$ of equation (2.1).

In its turn, from Lemma 2 and Theorem 1 it follows

Theorem 2. *If for function $\widetilde{M}(\varepsilon)$ equalling to right hand (2.6) there exists a number $\alpha \geq \beta > 0$ such that the inequalities (2.7) are true, then there exists $t(\varepsilon) \rightarrow \infty$ such that in $\mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$ there is one eigenvalue λ_ε of operator H_ε and it has asymptotics given by (2.3), (2.5), (2.6).*

If for the function $\widetilde{M}(\varepsilon)$ equalling to right hand of (2.6) there exists a number $\alpha > 0$ such that one of inequalities (2.8), (2.9) holds then there exists $t(\varepsilon) \rightarrow \infty$, such that in $\mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$ there is no eigenvalue of operator H_ε .

In particular, it follows from Theorem 2, that if the eigenvalue exists, it has asymptotics

$$\lambda_\varepsilon = -4 \exp \left\{ -\frac{4\pi}{\varepsilon \sum_{j=0}^{\infty} (-\varepsilon)^j \left\langle (\mathcal{L}_\varepsilon \circ \Delta^{-1})^j \mathcal{L}_\varepsilon[1] \right\rangle} - 2\mathbf{C} \right\}.$$

Remark 2. The formulation of Theorem 2 means that if one of its assumption takes place only on some subsequence $\varepsilon_j \rightarrow 0$, then the corresponding statement holds true on this subsequence.

3. PROOF OF THEOREM 1

We indicate by $\mathcal{B}^m(X, Y)$ (by $\mathcal{B}^m(X)$) the set of meromorphic operator-valued functions with values in $\mathcal{B}(X, Y)$ (in $\mathcal{B}(X)$). Let $\mathcal{B}(X, W_{2,loc}^2(\mathbb{R}^2))$ be a set of linear operators from Banach space X in $W_{2,loc}^2(\mathbb{R}^2)$ whose restriction on each bounded subset D belongs to $\mathcal{B}(X, W_2^2(D))$. Similarly, by $\mathcal{B}^h(X, W_{2,loc}^2(\mathbb{R}^2))$ (by $\mathcal{B}^m(X, W_{2,loc}^2(\mathbb{R}^2))$) we denote the set of operator-valued functions with values in $\mathcal{B}(X, W_{2,loc}^2(\mathbb{R}^2))$ such that for each bounded D they belong to $\mathcal{B}^h(X, W_2^2(D))$ (belong to $\mathcal{B}^m(X, W_2^2(D))$). Let $P_\varepsilon(k)$ be an operator defined by equality

$$P_\varepsilon(k)f = -\varepsilon \frac{(\ln k + \mathbf{C} - \ln 2) \langle B_\varepsilon(k)f \rangle B_\varepsilon(k) \mathcal{L}_\varepsilon[1]}{2\pi + \varepsilon(\ln k + \mathbf{C} - \ln 2) \langle B_\varepsilon(k) \mathcal{L}_\varepsilon[1] \rangle} + B_\varepsilon(k)f,$$

$$\mathcal{R}_\varepsilon(k) \stackrel{\text{def}}{=} A(k)P_\varepsilon(k).$$

Theorem 3. *For each $R > 0$ there exists $\varepsilon_0(R) > 0$ such that for $\varepsilon < \varepsilon_0$ and $k \in S_-(R)$*

1) $\mathcal{R}_\varepsilon(k) \in \mathcal{B}^m(L_2(\mathbb{R}^2; Q), W_{2,loc}^2(\mathbb{R}^2))$, and also, poles k_ε of operator $\mathcal{R}_\varepsilon(k)$ coincide with solutions of equation (2.1); if, in addition, $k \in \mathbb{C}_+$, then $\mathcal{R}_\varepsilon(k) \in \mathcal{B}^m(L_2(\mathbb{R}^2, Q); W_2^2(\mathbb{R}^2))$;

2) for each $f \in L_2(\mathbb{R}^2; Q)$ the function $u_\varepsilon = \mathcal{R}_\varepsilon(k)f$ is a solution of the equation

$$-H_\varepsilon u_\varepsilon = k^2 u_\varepsilon + f \quad \text{in } \mathbb{R}^2; \quad (3.1)$$

3) the residue of function u_ε at pole k_ε is determined by equality (2.4) up to multiplicative constant, moreover, this constant is nonzero if $\langle f \rangle \neq 0$.

Proof. By definition $A(k) \in \mathcal{B}^h(L_2(\mathbb{R}^2; Q), W_{2,loc}^2(\mathbb{R}^2))$ as $k \in S_-(R)$ for each R , moreover, $A(k) \in \mathcal{B}^h(L_2(\mathbb{R}^2; Q), W_2^2(\mathbb{R}^2))$ as $k \in \mathbb{C}_+$. Hence, taking into account the definition of $\mathcal{R}_\varepsilon(k)$ and Lemma 1, we get the validity of statement 1) of the theorem being proved.

Let us show the correctness of item 2). We seek the solution of equation (3.1) as

$$u_\varepsilon = A(k)g_\varepsilon, \quad (3.2)$$

where g_ε is a function to be found belonging to $L_2(\mathbb{R}^2; Q)$. Substituting (3.2) into (3.1) and bearing in mind that (3.2) is a solution of the equation

$$-H_0 u_\varepsilon = k^2 u_\varepsilon + g_\varepsilon \quad \text{in } \mathbb{R}^2,$$

we get that (3.2) is a solution of (3.1) in the case when

$$(I + \varepsilon T_\varepsilon(k))g_\varepsilon = f, \quad (3.3)$$

where

$$T_\varepsilon(k) = \mathcal{L}_\varepsilon A(k). \quad (3.4)$$

It follows from (3.4) and definition of $T_\varepsilon^{(0)}(k)$ and $A(k)$ that the operator $T_\varepsilon(k)$ acts in a following way:

$$T_\varepsilon(k)g = \frac{\langle g \rangle \mathcal{L}_\varepsilon[1]}{2\pi} (\ln k + \mathbf{C} - \ln 2) + T_\varepsilon^{(0)}(k)g. \quad (3.5)$$

Let $R > 0$ be an arbitrary number and ε satisfy to Lemma 1. Applying the operator $B_\varepsilon(k)$ to both hands of (3.3) and taking into account (3.5), we obtain that

$$\left(g_\varepsilon + \varepsilon \frac{\langle g_\varepsilon \rangle (\ln k + \mathbf{C} - \ln 2)}{2\pi} B_\varepsilon(k) \mathcal{L}_\varepsilon[1] \right) = B_\varepsilon(k)f. \quad (3.6)$$

Having integrated (3.6), we get that

$$\langle g_\varepsilon \rangle \left(1 + \varepsilon \frac{(\ln k + \mathbf{C} - \ln 2)}{2\pi} \langle B_\varepsilon(k) \mathcal{L}_\varepsilon[1] \rangle \right) = \langle B_\varepsilon(k)f \rangle. \quad (3.7)$$

Calculating $\langle g_\varepsilon \rangle$ by (3.7) and substituting this value in (3.6), we deduce that

$$g_\varepsilon = P_\varepsilon(k)f. \quad (3.8)$$

Assertions (3.2) and (3.8) imply the validity of statement 2). In its turn, the correctness 3) is yielded by 1), 2) and definition of $\mathcal{R}_\varepsilon(k)$. The proof is complete.

We denote by $R_\varepsilon(\lambda)$ the resolvent for operator H_ε .

Lemma 3. *The number of poles of resolvent $R_\varepsilon(\lambda)$, their orders and dimensions of residua at them are completely determined by functions from $L_2(\mathbb{R}^2; Q)$.*

Proof. Let F be an arbitrary function having finite support D . It is easy to see that

$$R_\varepsilon(\lambda)F = R_0(\lambda)F + \varepsilon R_\varepsilon(\lambda) \mathcal{L}_\varepsilon R_0(\lambda)F. \quad (3.9)$$

Since $R_0(\lambda)$ has no poles, and $\text{supp } (\mathcal{L}_\varepsilon R_0(\lambda)F) \subset \overline{Q}$, it follows from (3.9) the statement of lemma.

We will employ the symbol $\Sigma(H_\varepsilon)$ for the set of eigenvalues of operator H_ε .

Theorem 4. *Let $R > 0$ be an arbitrary number and $\varepsilon_0(R)$ satisfy to statement of Theorem 3, $\lambda = -k^2$. Then*

$$\mathcal{R}_\varepsilon(k)f = -R_\varepsilon(\lambda)f \quad (3.10)$$

for each $f \in L_2(\mathbb{R}^2; Q)$, $0 < \varepsilon < \varepsilon_0(R)$ and $k \in S^+(R)$ (or, equivalently, for each $\lambda \in S_+(R^2)$).

If $k_\varepsilon \in S_-(R)$ is a pole of operator $\mathcal{R}_\varepsilon(k)$ and $\text{Re } k_\varepsilon \leq 0$, then $\Sigma(H_\varepsilon) \cap S_+(R^2) = \emptyset$.

If $k_\varepsilon \in S^+(R)$ is a pole of operator $\mathcal{R}_\varepsilon(k)$, then $\Sigma(H_\varepsilon) \cap S_+(R^2) = \{\lambda_\varepsilon\}$, where λ_ε and the associated unique eigenfunction are given by equalities (2.3) and (2.4).

Proof. The function $\lambda = -k^2$ mapping $S^+(R)$ onto $S_+(R^2)$ in one-to-one manner, the correctness of equality (3.10) follows from statement 2) of Theorem 3 and the definition of resolvent. The correctness of other statement of the theorem being proved follows from Theorem 3 and Lemma 3.

Lemma 4. $\Sigma(H_\varepsilon) \subset \Pi_\varepsilon(-\varepsilon C(\mathcal{L}))$. If $\mathcal{L}_\varepsilon$ is a real operator then $\Sigma(H_\varepsilon) \subset [-\varepsilon C(\mathcal{L}), \infty)$.

Proof. Let $\lambda_\varepsilon \in \Sigma(H_\varepsilon)$, ϕ_ε be an associated eigenfunction normalized in $L_2(\mathbb{R}^2)$. Therefore

$$H_\varepsilon \phi_\varepsilon = \lambda_\varepsilon \phi_\varepsilon \quad \text{in } \mathbb{R}^2. \quad (3.11)$$

Since $H_\varepsilon U = H_0 U$ outside $\overline{Q}$, it is easy to show that $\phi_\varepsilon \in W_2^2(\mathbb{R}^2)$. Multiplying both hands of (3.11) by $\overline{\phi_\varepsilon}$ and integrating by parts we obtain the equality

$$\|\nabla \phi_\varepsilon\|_{\mathbb{R}^2}^2 - \varepsilon \langle \overline{\phi_\varepsilon} \mathcal{L}_\varepsilon \phi_\varepsilon \rangle = \lambda_\varepsilon. \quad (3.12)$$

Taking imaginary and real part of (3.12) and using an estimate $\|\mathcal{L}_\varepsilon \phi_\varepsilon\|_Q \leq C(\mathcal{L}) \|\phi_\varepsilon\|_{2,Q}$, we conclude that the lemma holds true.

Let $[a, b]_{\mathbb{R}}$ be a segment $[a, b]$ of real axis.

Lemma 5. For each $R > 0$ there exists $\varepsilon_0(R) > 0$ such that $[0, R]_{\mathbb{R}} \cap \Sigma(H_\varepsilon) = \emptyset$ as $\varepsilon < \varepsilon_0$.

Proof. We prove the lemma by reductio ad absurdum. It is easy to see that without loss of generality it is sufficient to consider the situation $[0, R]_{\mathbb{R}} \cap \Sigma(H_\varepsilon) = \lambda_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} \lambda_*$ in the cases when all $\lambda_\varepsilon \neq 0$, and, conversely, when all $\lambda_\varepsilon = 0$. Clear, in the latter case $\lambda_* = 0$.

We start from first case. Since $H_\varepsilon U = H_0 U$ outside $\overline{Q}$, then for the associated eigenfunction ϕ_ε the equality $\phi_\varepsilon = 0$ holds as $|x| = T$ for each T sufficiently large (see., for instance, [14, § 3.10]). Hence, λ_ε and ϕ_ε are eigenelements of the boundary value problem

$$H_\varepsilon \phi_\varepsilon = \lambda_\varepsilon \phi_\varepsilon \quad \text{as } |x| < T, \quad \phi_\varepsilon = 0 \quad \text{as } |x| = T. \quad (3.13)$$

Since (3.13) is a regular perturbation of the limit problem

$$H_0 \phi_* = \lambda_* \phi_* \quad \text{as } |x| < T, \quad \phi_* = 0 \quad \text{as } |x| = T, \quad (3.14)$$

it follows that having chosen T in such a way that λ_* not to be an eigenvalue of (3.14), we get that ϕ_ε can not satisfy to (3.13).

Let us proceed to the second case $\lambda_\varepsilon = 0$. In an obvious way from (3.11) and properties of $\mathcal{L}_\varepsilon$ it follows the estimate

$$\begin{aligned} \|\Delta \phi_\varepsilon\|_{\mathbb{R}^2} &\leq \varepsilon C \|\phi_\varepsilon\|_{2,Q}, \\ \|\nabla \phi_\varepsilon\|_{\mathbb{R}^2}^2 &\leq \varepsilon C \|\phi_\varepsilon\|_Q \|\phi_\varepsilon\|_{2,Q}. \end{aligned} \quad (3.15)$$

Without loss of generality we assume that the function ϕ_ε is normalized in $W_2^2(Q)$. Then the estimates (3.15) imply that on Q

$$\phi_\varepsilon = |Q|^{-1/2} + \psi_\varepsilon, \quad (3.16)$$

$$\|\psi_\varepsilon\|_{1,Q} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (3.17)$$

By localness of $\mathcal{L}_\varepsilon$ and (3.16) outside Q the function ϕ_ε being a solution of boundary value problem

$$\Delta\phi_\varepsilon = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{Q} \quad \phi_\varepsilon = |Q|^{-1/2} + \psi_\varepsilon \quad \text{on } \partial Q,$$

it follows from (3.17) that ϕ_ε does not tends to zero at infinity. Therefore, $\phi_\varepsilon \notin L_2(\mathbb{R}^2)$. The contradiction obtained completes the proof of lemma.

One can easily see that Theorem 1 is a direct implication of Theorem 4 and Lemmas 4 and 5.

4. EXAMPLES

Example 1. Let $\mathcal{L}_\varepsilon[g] = Vg$, where $V \in C_0^\infty(Q)$. It follows from (2.6) that for this case

$$\widetilde{M}(\varepsilon) = \frac{2\pi}{\varepsilon m(\varepsilon)} + \mathbf{C} - \ln 2, \quad (4.1)$$

where

$$m(\varepsilon) = \langle V \rangle + \sum_{j=1}^3 (-\varepsilon)^j \langle (V \Delta^{-1})^j V \rangle + O(\varepsilon^4). \quad (4.2)$$

We note, that by integrating by parts it is easy to show the validity of equalities:

$$\begin{aligned} \langle U \Delta^{-1} U \rangle &= - \langle (\nabla (\Delta^{-1} U))^2 \rangle, \quad \langle (U \Delta^{-1})^2 U \rangle = \langle U (\Delta^{-1} U)^2 \rangle, \\ \langle (U \Delta^{-1})^3 U \rangle &= \langle (U \Delta^{-1} U) \Delta^{-1} (U \Delta^{-1} U) \rangle, \quad \text{if } \langle U \rangle = 0. \end{aligned} \quad (4.3)$$

The assertions (4.1)–(4.3) yield

$$\widetilde{M}(\varepsilon) = \frac{2\pi}{\varepsilon \langle V \rangle} + \frac{2\pi \langle V \Delta^{-1} V \rangle}{\langle V \rangle^2} + \mathbf{C} - \ln 2 + O(\varepsilon), \quad \text{if } \langle V \rangle \neq 0, \quad (4.4)$$

$$\begin{aligned} \widetilde{M}(\varepsilon) &= \frac{2\pi}{\varepsilon^2 \langle (\nabla (\Delta^{-1} V))^2 \rangle} - \frac{2\pi \langle V (\Delta^{-1} V)^2 \rangle}{\varepsilon \langle (\nabla (\Delta^{-1} V))^2 \rangle^2} \\ &\quad + \frac{2\pi \langle (V \Delta^{-1} V) \Delta^{-1} (V \Delta^{-1} V) \rangle}{\langle (\nabla (\Delta^{-1} V))^2 \rangle^2} + \frac{2\pi \langle V (\Delta^{-1} V)^2 \rangle^2}{\langle (\nabla (\Delta^{-1} V))^2 \rangle^3} \\ &\quad + \mathbf{C} - \ln 2 + O(\varepsilon), \quad \text{if } \langle V \rangle = 0. \end{aligned} \quad (4.5)$$

First we consider classical situation $\text{Im } V = 0$. It follows from (4.4) and Theorem 2 that if $\langle V \rangle < 0$, then there is no eigenvalue converging to zero, and if $\langle V \rangle > 0$, then

$$\lambda_\varepsilon = -\varkappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon \langle V \rangle} \right\}, \quad (4.6)$$

$$\varkappa_\varepsilon = 2 \exp \left\{ -\frac{2\pi \langle V \Delta^{-1} V \rangle}{\langle V \rangle^2} - \mathbf{C} \right\} + O(\varepsilon).$$

By analogy, from (4.5) and Theorem 2 it follows that if $\langle V \rangle = 0$, then the eigenvalue exists and has the asymptotics

$$\lambda_\varepsilon = -\varkappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon^2 \|\nabla(\Delta^{-1}V)\|_{\mathbb{R}^2}^2} + \frac{4\pi \langle V(\Delta^{-1}V)^2 \rangle}{\varepsilon \|\nabla(\Delta^{-1}V)\|_{\mathbb{R}^2}^4} \right\}, \quad (4.7)$$

$$\varkappa_\varepsilon = 2 \exp \left\{ -\frac{2\pi \langle (V\Delta^{-1}V)\Delta^{-1}(V\Delta^{-1}V) \rangle}{\|\nabla(\Delta^{-1}V)\|_{\mathbb{R}^2}^4} - \frac{2\pi \langle V(\Delta^{-1}V)^2 \rangle^2}{\|\nabla(\Delta^{-1}V)\|_{\mathbb{R}^2}^6} - \mathbf{C} \right\} + O(\varepsilon).$$

It should be noted that formulas (4.6), (4.7) are well-known (see [1], [2]).

Now let us consider complex potential V . It follows from Theorem 2 and (4.4) that if $\text{Im } \langle V \rangle \neq 0$, then for all sufficiently small ε there is no eigenvalue outside $\mathbb{C} \setminus \Pi_\varepsilon(t(\varepsilon))$ (i.e., there is no small eigenvalue). Observe, this situation differs noticeably from one-dimensional case in [11], where the sufficient condition for the existence of the eigenvalue converging to zero was $\text{Re } \langle V \rangle > 0$.

Let us show that for complex potential even the condition $\langle V \rangle > 0$ is insufficient for existence of eigenvalue. Let $V = v + ia\Delta v$, where $v \in C_0^\infty(Q)$ is a real function, $\langle v \rangle > 0$, and a is an arbitrary real constant unknown yet. It is easy to see that in this case

$$\langle V \rangle = \langle v \rangle > 0, \quad \langle V\Delta^{-1}V \rangle = (1 - a^2) \langle v\Delta^{-1}v \rangle + i2a \|v\|_{\mathbb{R}^2}^2.$$

Then it follows from (4.4) that

$$\begin{aligned} \text{Re } \widetilde{M}(\varepsilon) &= \frac{2\pi}{\varepsilon \langle v \rangle} + \frac{2\pi(1 - a^2) \langle v\Delta^{-1}v \rangle}{\langle v \rangle^2} + \mathbf{C} - \ln 2 + O(\varepsilon), \\ \text{Im } \widetilde{M}(\varepsilon) &= \frac{4\pi a \|v\|_{\mathbb{R}^2}^2}{\langle v \rangle^2} + O(\varepsilon). \end{aligned} \quad (4.8)$$

It follows from (4.8) and Theorem 2 that for

$$|a| > \frac{\langle v \rangle^2}{8 \|v\|^2}$$

there is no small eigenvalue, and for

$$|a| < \frac{\langle v \rangle^2}{8 \|v\|^2}$$

such eigenvalue exists and has asymptotics (4.6), where

$$\begin{aligned} \varkappa_\varepsilon &= 2 \exp \left\{ \frac{2\pi(a^2 - 1) \langle v\Delta^{-1}v \rangle}{\langle v \rangle^2} - \mathbf{C} \right\} \\ &\times \left(\cos \left(\frac{4a\pi \|v\|^2}{\langle v \rangle^2} \right) - i \sin \left(\frac{4a\pi \|v\|^2}{\langle v \rangle^2} \right) \right) + O(\varepsilon). \end{aligned} \quad (4.9)$$

In particular, it follows from (4.6) and (4.9) that for

$$\frac{\langle v \rangle^2}{16 \|v\|^2} < |a| < \frac{\langle v \rangle^2}{8 \|v\|^2}$$

the eigenvalue lies in the right complex half-plane (in contrast to the case of real potential, for that eigenvalue can lie only in the negative real semiaxis).

Example 2. Let $\mathcal{L}_\varepsilon[g] = V_\varepsilon g$, where $V_\varepsilon = V + \varepsilon V_1$, and V, V_1 are real function having supports in Q . Then by (2.6), (4.3) and by equalities checked easily

$$\begin{aligned} \langle V_1 \Delta^{-1} V \rangle &= \langle V \Delta^{-1} V_1 \rangle, & \left\langle (V \Delta^{-1})^2 V_1 \right\rangle &= \langle V (\Delta^{-1} V) (\Delta^{-1} V_1) \rangle, \\ \langle V \Delta^{-1} (V_1 \Delta^{-1} V) \rangle &= \left\langle V_1 (\Delta^{-1} V)^2 \right\rangle & \text{as } \langle V \rangle = 0 \end{aligned}$$

$\widetilde{M}(\varepsilon)$ has the form (4.1), where

$$m(\varepsilon) = \langle V \rangle + \varepsilon (\langle V_1 \rangle - \langle V \Delta^{-1} V \rangle) + O(\varepsilon^2), \quad \text{if } \langle V \rangle \neq 0, \quad (4.10)$$

and

$$\begin{aligned} \frac{m(\varepsilon)}{\varepsilon} &= \|\nabla (\Delta^{-1} V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle - \varepsilon \left\{ 2 \langle V_1 \Delta^{-1} V \rangle - \left\langle V (\Delta^{-1} V)^2 \right\rangle \right\} \\ &\quad - \varepsilon^2 m_2 + O(\varepsilon^3), \quad \text{if } \langle V \rangle = 0, \end{aligned} \quad (4.11)$$

where

$$\begin{aligned} m_2 &= \langle V_1 \Delta^{-1} V_1 \rangle + \langle (V \Delta^{-1} V) \Delta^{-1} (V \Delta^{-1} V) \rangle \\ &\quad - \langle V (\Delta^{-1} V) (\Delta^{-1} V_1) \rangle - \left\langle V_1 (\Delta^{-1} V)^2 \right\rangle - \langle V_1 \Delta^{-1} (V \Delta^{-1} V) \rangle. \end{aligned} \quad (4.12)$$

It follows from (4.1), (4.10), (4.11) that

$$\widetilde{M}(\varepsilon) = \frac{2\pi}{\varepsilon \langle V \rangle} + \frac{2\pi (\langle V \Delta^{-1} V \rangle - \langle V_1 \rangle)}{\langle V \rangle^2} + \mathbf{C} - \ln 2 + O(\varepsilon), \quad \text{if } \langle V \rangle \neq 0, \quad (4.13)$$

and

$$\begin{aligned} \widetilde{M}(\varepsilon) &= \frac{2\pi}{\varepsilon^2 (\|\nabla (\Delta^{-1} V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)} + \frac{2\pi \left(2 \langle V_1 \Delta^{-1} V \rangle - \left\langle V (\Delta^{-1} V)^2 \right\rangle \right)}{\varepsilon (\|\nabla (\Delta^{-1} V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^2} \\ &\quad + \frac{2\pi \left(2 \langle V_1 \Delta^{-1} V \rangle - \left\langle V (\Delta^{-1} V)^2 \right\rangle \right)^2}{(\|\nabla (\Delta^{-1} V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^3} + \frac{2\pi m_2}{(\|\nabla (\Delta^{-1} V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^2} \\ &\quad + \mathbf{C} - \ln 2 + O(\varepsilon), \quad \text{if } \langle V \rangle = 0. \end{aligned} \quad (4.14)$$

It follows from (4.13), (4.14) and Theorem 2 that the condition $\langle V_\varepsilon \rangle \geq 0$ is sufficient for existence of eigenvalue. If $\langle V_\varepsilon \rangle > 0$ "in principle" (i.e., $\langle V \rangle > 0$), then by (4.13) it has asymptotics (4.6), where

$$\varkappa_\varepsilon = 2 \exp \left\{ -\frac{2\pi (\langle V \Delta^{-1} V \rangle - \langle V_1 \rangle)}{\langle V \rangle^2} - \mathbf{C} \right\} + O(\varepsilon).$$

If $\langle V_\varepsilon \rangle > 0$ in small (i.e., $\langle V \rangle = 0$, but $\langle V_1 \rangle > 0$), then due to (4.14) the eigenvalue has asymptotics

$$\begin{aligned}\lambda_\varepsilon &= -\kappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon^2 (\langle V_1 \rangle + \|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2)} \right. \\ &\quad \left. - \frac{4\pi \left(2\langle V_1 \Delta^{-1}V \rangle - \langle V (\Delta^{-1}V)^2 \rangle \right)}{\varepsilon (\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^2} \right\}, \\ \kappa_\varepsilon &= 2 \exp \left\{ -\frac{2\pi \left(2\langle V_1 \Delta^{-1}V \rangle - \langle V (\Delta^{-1}V)^2 \rangle \right)^2}{(\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^3} \right. \\ &\quad \left. - \frac{2\pi m_2}{(\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^2} - \mathbf{C} \right\} + O(\varepsilon),\end{aligned}$$

where constant m_2 is defined by (4.12). And, finally, if $\langle V_\varepsilon \rangle = 0$, then in view of (4.14) and (4.3) the eigenvalue has asymptotics

$$\begin{aligned}\lambda_\varepsilon &= -\kappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon^2 \|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2} - \frac{4\pi \left(2\langle V_1 \Delta^{-1}V \rangle + \langle V (\Delta^{-1}V)^2 \rangle \right)}{\varepsilon \|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^4} \right\}, \\ \kappa_\varepsilon &= 2 \exp \left\{ -\frac{2\pi \left(2\langle V_1 \Delta^{-1}V \rangle + \langle V (\Delta^{-1}V)^2 \rangle \right)^2}{\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^6} - \frac{2\pi m^{(2)}}{\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^4} - \mathbf{C} \right\} + O(\varepsilon),\end{aligned}$$

where

$$\begin{aligned}m^{(2)} &= -\|\nabla (\Delta^{-1}V_1)\|^2 + \langle (V \Delta^{-1}V) \Delta^{-1} (V \Delta^{-1}V) \rangle \\ &\quad - \langle V (\Delta^{-1}V) (\Delta^{-1}V_1) \rangle - \langle V_1 (\Delta^{-1}V)^2 \rangle - \langle V_1 \Delta^{-1} (V \Delta^{-1}V) \rangle.\end{aligned}$$

However, in contrast to classic case (real $V_\varepsilon = V$) the condition $\langle V_\varepsilon \rangle < 0$ is insufficient for absence of eigenvalue. Indeed, let v be a real function, $V = \Delta v$, $V_1 = av$, $\langle v \rangle > 0$ and a is a some arbitrary constant. Then by (4.14)

$$\begin{aligned}\widetilde{M}(\varepsilon) &= \frac{2\pi}{\varepsilon^2 (\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)} + \frac{2\pi (2a \|\nabla v\|_{\mathbb{R}^2}^2 - \langle v^2 \Delta v \rangle)}{\varepsilon (\|\nabla (\Delta^{-1}V)\|_{\mathbb{R}^2}^2 + \langle V_1 \rangle)^2} \\ &\quad + \frac{2\pi (2a \|\nabla v\|_{\mathbb{R}^2}^2 - \langle v^2 \Delta v \rangle)^2}{(\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)^3} + \frac{2\pi \widetilde{m}_2}{(\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)^2} \\ &\quad + \mathbf{C} - \ln 2 + O(\varepsilon),\end{aligned}\tag{4.15}$$

where

$$\begin{aligned}\widetilde{m}_2 &= a^2 \langle v \Delta^{-1}v \rangle + \langle (v \Delta v) \Delta^{-1} (v \Delta v) \rangle \\ &\quad - a \langle v (\Delta^{-1}v) \Delta v \rangle - a \langle v^3 \rangle - a \langle v \Delta^{-1} (v \Delta v) \rangle.\end{aligned}\tag{4.16}$$

It follows from (4.15) and Theorem 2 that for

$$-\frac{\|\nabla v\|^2}{\langle v \rangle} < a < 0$$

the eigenvalue exists and has asymptotics

$$\lambda_\varepsilon = -\kappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon^2 (\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)} - \frac{4\pi (2a \|\nabla v\|_{\mathbb{R}^2}^2 - \langle v^2 \Delta v \rangle)}{\varepsilon (\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)^2} \right\},$$

$$\kappa_\varepsilon = 2 \exp \left\{ -\frac{2\pi (2a \|\nabla v\|_{\mathbb{R}^2}^2 - \langle v^2 \Delta v \rangle)^2}{(\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)^3} - \frac{2\pi \tilde{m}_2}{(\|\nabla v\|_{\mathbb{R}^2}^2 + a \langle v \rangle)^2} - \mathbf{C} \right\} + O(\varepsilon),$$

where $\tilde{m}_2$ is defined in (4.16). At the same time, clearly, $\langle V_\varepsilon \rangle = a \langle v \rangle < 0$ for the indicated values of a . Similar effect was observed in [6].

Example 3. Let

$$\mathcal{L}_\varepsilon[g] = \chi(Q) \langle \rho_\varepsilon g \rangle,$$

where $\chi(Q)$ is a characteristic function of Q , (i.e., the function equalling to one as $x \in Q$ and vanishing for other x), and a function ρ_ε be continuous in $\overline{Q}$ and extended by zero outside $\overline{Q}$.

Since

$$\mathcal{L}_\varepsilon[1] = \langle \rho_\varepsilon \rangle, \tag{4.17}$$

it follows from Corollary of Theorem 1 that there is no small eigenvalue if $\langle \rho_\varepsilon \rangle = 0$.

From definition of $\mathcal{L}_\varepsilon$ it follows that

$$(\mathcal{L}_\varepsilon \circ \Delta^{-1})^n \mathcal{L}_\varepsilon[1] = \langle \rho_\varepsilon \Delta^{-1} \chi \rangle^n \chi, \quad n \geq 1. \tag{4.18}$$

Substituting (4.17), (4.18) into (2.6) we get that

$$\widetilde{M}(\varepsilon) = \frac{2\pi}{\varepsilon \langle \rho_\varepsilon \rangle |Q|} (1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle) + \mathbf{C} - \ln 2. \tag{4.19}$$

We remind, that in view of Theorem 2 the sufficient condition for existence of small eigenvalue is an existence of $\alpha > 0$ such that the inequalities (2.7) hold, due to (4.19) they take the form

$$\operatorname{Re} \frac{1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle}{\varepsilon \langle \rho_\varepsilon \rangle} > \alpha \varepsilon^\alpha + \frac{|Q| (\ln 2 - \mathbf{C})}{2\pi},$$

$$\left| \operatorname{Im} \frac{1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle}{\varepsilon \langle \rho_\varepsilon \rangle} \right| < \frac{|Q|}{4} - \alpha \varepsilon^\alpha. \tag{4.20}$$

The first of these inequalities holds true, if quantities $\langle \rho_\varepsilon \Delta^{-1} \chi \rangle$ and $\langle \rho_\varepsilon \rangle$ are bounded by module as $\varepsilon \rightarrow 0$ and $\operatorname{Re} \langle \rho_\varepsilon \rangle > 0$, and second relationship, clear, take place for

real ρ_ε . Under fulfilment of condition (4.20) due to Theorem 2 the eigenvalue is of the form

$$\lambda_\varepsilon = -4 \exp \left\{ -\frac{4\pi (1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle)}{\varepsilon |Q| \langle \rho_\varepsilon \rangle} - 2\mathbf{C} + O(\varepsilon^\infty) \right\}. \quad (4.21)$$

By analogy, due to Theorem 2 the sufficient condition of absence small eigenvalue is an existence of $\alpha > 0$ such that the inequalities (2.8) or (2.9) hold true, those due to (4.19) are of the form

$$\operatorname{Re} \frac{1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle}{\varepsilon \langle \rho_\varepsilon \rangle} < -\alpha \varepsilon^\alpha + \frac{|Q| (\ln 2 - \mathbf{C})}{2\pi}$$

or

$$\left| \operatorname{Im} \frac{1 + \varepsilon \langle \rho_\varepsilon \Delta^{-1} \chi \rangle}{\varepsilon \langle \rho_\varepsilon \rangle} \right| > \frac{|Q|}{4} + \alpha \varepsilon^\alpha.$$

If quantities $\langle \rho_\varepsilon \Delta^{-1} \chi \rangle$ and $\langle \rho_\varepsilon \rangle$ are bounded by module as $\varepsilon \rightarrow 0$ and $\operatorname{Re} \langle \rho_\varepsilon \rangle < 0$, then first of these inequalities holds a fortiori, and second inequality, obviously, take place for $|\operatorname{Im} \langle \rho_\varepsilon \rangle| > c > 0$.

Remark 3. Defining concretely ρ_ε (like it was done in Examples 1 and 2), by simple relationships (4.20) and (4.21) it is easy to get more explicit formulas for eigenvalues (like in Examples 1, 2). In particular, from (4.20) and (4.21) it follows that if $\rho_\varepsilon \equiv 1$ in Q , then

$$\begin{aligned} \lambda_\varepsilon &= -\varkappa_\varepsilon^2 \exp \left\{ -\frac{4\pi}{\varepsilon |Q|^2} \right\}, \\ \varkappa_\varepsilon &= 2 \exp \left\{ -\frac{2\pi \langle \Delta^{-1} \chi \rangle}{|Q|^2} - \mathbf{C} \right\} + O(\varepsilon^\infty). \end{aligned}$$

On the other hand, for the case $\mathcal{L}_\varepsilon = V_\varepsilon$ considered in Examples 1 and 2 under fulfilment of inequalities (2.7) from Theorem 2 and (2.6) we deduce the following "general" analog of formula (4.21) for eigenvalues:

$$\lambda_\varepsilon = -4 \exp \left\{ -\frac{4\pi}{\varepsilon \sum_{j=0}^{\infty} (-\varepsilon)^j \left\langle (V_\varepsilon \circ \Delta^{-1})^j V_\varepsilon \right\rangle} - 2\mathbf{C} + O(\varepsilon^\infty) \right\},$$

that is, however, is not such constructive like (4.21) and formulas obtained in Examples 1 and 2.

Example 4. Let

$$\mathcal{L}_\varepsilon g = \sum_{i,j=1}^2 \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial}{\partial x_j} g \right) + \sum_{i=1}^2 \frac{\partial}{\partial x_i} (a_i g) + \mathcal{L}^\varepsilon g, \quad (4.22)$$

where $a_{ij}, a_i, \in C_0^\infty(Q)$,

$$\mathcal{L}^\varepsilon \stackrel{\text{def}}{=} V_\varepsilon \quad \text{or} \quad \mathcal{L}^\varepsilon[g] \stackrel{\text{def}}{=} \chi(Q) \langle \rho_\varepsilon g \rangle.$$

Since $\mathcal{L}_\varepsilon[1] = \mathcal{L}^\varepsilon[1]$ and $\langle \mathcal{L}_\varepsilon g \rangle = \langle \mathcal{L}^\varepsilon g \rangle$, by (2.6) the function $\widetilde{M}(\varepsilon)$, corresponding to the operator defined by (4.22), coincides with the function $\widetilde{M}(\varepsilon)$, corresponding to the operator $\mathcal{L}_\varepsilon = \mathcal{L}^\varepsilon$. Therefore, all the results established in Examples 1–3 (for operator $\mathcal{L}_\varepsilon = V_\varepsilon$ and operator defined by equality $\mathcal{L}^\varepsilon[g] = \chi(Q) \langle \rho_\varepsilon g \rangle$), are carried over with no changes to the case of operator defined by equality (4.22).

Acknowledgments. The work has done under support of RFBR grants (02-01-00693, 00-15-96038) and Ministry of Education of Russia (E00-1.0-53).

References

- [1] *L. D. Landau, E. M. Lifshits.* Theoretical physics. Vol. 3. Quantum mechanics. Non-relativistic theory. M.: Nauka, 1974. (in Russian)
- [2] *B. Simon.* Ann. Phys. 1976. V. 97. P. 279–288.
- [3] *M. Klaus.* Ann. Phys. 1977. V. 108. P. 288–300.
- [4] *R. Blankenbecler, M.L. Goldberger, B. Simon.* Ann. Phys. 1977. V. 108. P. 69–78.
- [5] *M. Klaus, B. Simon.* Ann. Phys. 1980. V. 130. P. 251–281.
- [6] *F. Bentosela, R.M. Cavalcanti, P.Exner, V.A. Zagrebanov.* J. Phys. A. 1999. V. 32. P. 3029–3039.
- [7] *D.E. Pelinovsky, C. Sulem.* Theor. Math. Phys. 2000. V. 122. P. 98–106.
- [8] *D.E. Pelinovsky, C. Sulem.* Commun. Math. Phys. 2000. V. 208. P. 713–760.
- [9] *P. Zhevandrov, A. Merzon.* Trans. AMS (accepted).
- [10] *P. Zhevandrov, A. Merzon.* Proc. 3-rd Intern. ISAAC Congress. Berlin. 2001 (accepted).
- [11] *R. R. Gadyl'shin.* Theor. Math. Phys. 2002. V. 132. P. 976–982.
- [12] *I. S. Gradshteyn, I. M. Ryzhik.* Table of integrals, series, and products. New York - London - Toronto: Academic Press, 1980.
- [13] *H. Bateman, A. Erdélyi.* Higher transcendental functions. Vol. 2. M.: "Nauka", 1974.
- [14] *D. Colton, R. Kress.* Integral equation methods in scattering theory. Pure and Applied Mathematics. A Wiley-Interscience Publication. New York etc.: John Wiley, 1983.